IN A NUTSHELL

The knowledge platform for general practitioners

Wells' Criteria for Pulmonary Embolism

The Wells' Criteria for pulmonary embolism can be used to estimate the probability of pulmonary embolism based on the history and clinical criteria. This results in a recommendation for further diagnostics depending on the pretest probability.

Clinical signs and symptoms of deep vein thrombosis (leg swelling or compressive tenderness along the deep veins) \bigcirc No \bigcirc Yes (+3) Pulmonary embolism as likely or more likely than an alternative diagnosis \bigcirc No \bigcirc Yes (+3) Hearth rate > 100 bpm \bigcirc No \bigcirc Yes (+1.5) Immobilization (? 3 days) or surgery in the past 4 weeks \bigcirc No \bigcirc Yes (+1.5) Previously diagnosed deep vein thrombosis or pulmonary embolism \bigcirc No \bigcirc Yes (+1.5) Hemoptysis \bigcirc No \bigcirc Yes (+1) Malignancy (existing therapy, palliative therapy, or treatment discontinuation within the last 6 months) \bigcirc No \bigcirc Yes (+1)

Wells' Criteria for Pulmonary Embolism:

Classification according van Belle et al. [2]

Recommendation:

Interpretation

1

Original classification according to Wells et Classification according to van Belle et al. Score al. [1] [2] Low probability of pulmonary embolism Prevalence: 1.2 % Pulmonary embolism unlikely Recommandation: perform D-dimer test Prevalence: 12.1 % ?1 - D-dimer negative: pulmonary embolism Recommendation: perform D-dimer test excluded - If D-dimer is negative, pulmonary embolism - D-dimer positive: imaging to rule out can be excluded. pulmonary embolism - If D-dimer is positive, CT angiography 2 should be performed to rule out pulmonary Moderate probability of pulmonary embolism. 3 embolism 4 Prevalence: 16.2 % Recommendation: imaging to rule out 5 pulmonary embolism 6 Pulmonary embolism likely Prevalence: 37.1 % High probability of pulmonary embolism Recommendation: CT angiography should be Prevalence: 37.5 % ?7 performed to rule out pulmonary embolism. Recommendation: imaging to rule out pulmonary embolism

References

- Wells PS, Anderson DR, Rodger M et al. Excluding pulmonary embolism at the bedside without diagnostic imaging: management of patients with suspected pulmonary embolism presenting to the emergency department by using a simple clinical model and d-dimer. Ann Intern Med 2001; 135: 98-107
- 2. van Belle A, Büller HR, Huisman MV et al. Effectiveness of managing suspected pulmonary embolism using an algorithm combining clinical probability, D-dimer testing, and computed tomography. JAMA 2006; 295: 172-179